

Depressive Symptoms and Quality of Life in Adolescents With Type 2 Diabetes

Baseline data from the TODAY study

BARBARA J. ANDERSON, PHD¹
SHARON EDELSTEIN, SCM²
NATALIE WALDERS ABRAMSON, PHD³
LORRAINE E. LEVITT KATZ, MD⁴
PATRICE M. YASUDA, PHD⁵
SYLVIA J. LAVIETES, MS⁶

PAULA M. TRIEF, PHD⁷
SHERIDA E. TOLLEFSEN, MD⁸
SIRIPOOM VUDHIPOOM MCKAY, MD⁹
PATRICIA KRINGAS, RN, MA, CDE¹⁰
TERRI LYNN CASEY, BSN, RN, CDE¹¹
MARSHA D. MARCUS, PHD¹²

RESEARCH DESIGN AND METHODS

Participants

The TODAY study design and participant characteristics have been reported (3). The protocol was approved by an External Evaluation Committee convened by the National Institute of Diabetes and Digestive and Kidney Diseases and by institutional review boards of each participating institution. Participants provided informed consent or assent.

Procedures

Eligible youth and an adult caregiver were recruited at 15 TODAY clinical centers (Supplementary Data). Relevant to the current report, youth were ineligible if diagnosed with major depression or if taking psychotropic medications. Before baseline, youth successfully completed a run-in phase lasting 8–24 weeks, during which youth had to establish blood glucose $HbA_{1c} < 8\%$ on metformin alone. Of the 927 youth who entered run-in, 704 (76%) successfully completed run-in and provided baseline data; 55 youth (5.9%) were not able to maintain glycemic control on metformin alone. Average HbA_{1c} at baseline was 5.9% (5.5–6.5). Detailed clinical characteristics of the TODAY cohort at baseline have been reported (3).

Depressive symptoms. The Children's Depression Inventory (CDI) (4), assessing depressive symptoms, was completed by participants aged ≤ 15 years. Items are combined into a total depressive symptom score. A cutoff score ≥ 13 identifies clinically significant levels of depressive symptoms (4).

The Beck Depression Inventory II (BDI-II) (5) was completed by participants aged ≥ 16 years. Items are combined into a total score, with a cutoff score ≥ 14 to identify a clinically significant level of depressive symptoms. Scores ≥ 29 are suggestive of "severe" depressive symptoms.

Quality of life. Pediatric Quality of Life (PedsQL) (6) measures the youth's perception of quality of life. The 23 items are scored using a five-point Likert scale; a total score is the average of all items.

OBJECTIVE—The study objective was to examine the prevalence of depressive symptoms and relationships to quality of life and demographics in the Treatment Options for Type 2 Diabetes in Adolescents and Youth (TODAY) study's large, ethnically diverse youth with type 2 diabetes.

RESEARCH DESIGN AND METHODS—A total of 704 youth with type 2 diabetes < 2 years' duration, aged 10–17 years, and BMI ≥ 85 th percentile completed depressive symptoms and quality of life measures.

RESULTS—Some 14.8% reported clinically significant depressive symptoms, and older girls had significantly higher rates than older boys.

CONCLUSIONS—Rates of significant depressive symptoms were similar to those of healthy adolescents and lower than those of teens with type 1 diabetes. Elevated depressive symptoms, particularly in older girls, suggest clinicians assess vulnerability.

Diabetes Care 34:2205–2207, 2011

Little is known about physiologic and psychosocial correlates of type 2 diabetes in adolescents, whereas data on adults with type 2 diabetes are available, with rates of depressive symptoms approximately twice as high as for adults without diabetes (1), underscoring the importance of examining depressive symptoms in adolescents with type 2 diabetes. Depression in adults is also

associated with significant psychosocial and physical complications (2). To date, we have not had access to large samples of patients needed to study depression systematically in youth with type 2 diabetes. Treatment Options for Type 2 Diabetes in Adolescents and Youth (TODAY) offers unique data concerning the psychosocial functioning of youth with type 2 diabetes.

From the ¹Department of Pediatrics, Section of Psychology, Baylor College of Medicine, Houston, Texas; the ²George Washington University Biostatistics Center, Rockville, Maryland; the ³Department of Psychiatry and Behavioral Sciences, The Children's Hospital, University of Colorado Denver School of Medicine, Aurora, Colorado; the ⁴Division of Endocrinology, Children's Hospital of Philadelphia/University of Pennsylvania School of Medicine, Philadelphia, Pennsylvania; the ⁵Center for Endocrinology, Diabetes & Metabolism, Children's Hospital Los Angeles, Los Angeles, California; the ⁶Department of Pediatric Diabetes, Yale University, New Haven, Connecticut; the ⁷Departments of Psychiatry and Medicine, SUNY Upstate Medical University, Syracuse, New York; the ⁸Department of Pediatrics, Saint Louis University Health Sciences Center, St. Louis, Missouri; the ⁹Department of Pediatric Endocrinology and Metabolism, Baylor College of Medicine, Houston, Texas; the ¹⁰Naomi Berrie Diabetes Center, Columbia University Medical Center, New York, New York; ¹¹Pediatric Diabetes, Endocrinology and Metabolism, University Hospitals Case Medical Center, Cleveland, Ohio; and the ¹²Western Psychiatric Institute and Clinic, University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania.

Corresponding author: Barbara J. Anderson, bja@bcm.edu.

Received 3 March 2011 and accepted 1 July 2011.

DOI: 10.2337/dc11-0431. Clinical trial reg. no. NCT00081328, clinicaltrials.gov.

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc11-0431/-/DC1>.

A list of the members of the TODAY Study Group can be found in the Supplementary Data online.

© 2011 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Statistical methods

Descriptive statistics at randomization are reported as means, SDs, or percents above cutoffs for serious level of depressive symptoms. Analysis of variance or *t* tests were used to compare mean depressive symptoms and χ^2 tests for percent above predefined cut points, by sex and race/ethnic group. Pearson correlation coefficients were used to compare the relationship between depressive symptoms and quality of life.

RESULTS—The demographic characteristics of the 704 youth randomized in TODAY have been reported (3). Briefly, girls comprised 65% of the sample, and 61% were aged 12–15 years. More than 80% reported minority race/ethnicity. Participants were overweight or obese, with 87% having a BMI >95th percentile for age and sex. Socioeconomic status was low, with >25% from homes in which neither parent had a high school diploma.

As shown in Table 1, 14.8% of youth reported depressive symptoms above the threshold for clinical significance. No differences in depressive symptom scores were seen across race/ethnicity; however, there were differences by sex (Table 1). BDI was significantly higher in older girls than in older boys, as was the percent scoring at or above the cutoff score for BDI and for CDI and BDI combined. No differences were found in the PedsQL or CDI for either race/ethnicity or sex. Both CDI and BDI were negatively correlated with PedsQL, -0.59 for the CDI and -0.63 for the BDI (both $P < 0.001$).

CONCLUSIONS—In the TODAY cohort, ~15% scored at or above the cutoff for clinically significant depressive symptoms, a rate similar to that of adolescents without type 2 diabetes (7). Depressive symptoms were not assessed before run-in. Thus, although it is not possible to rule out that completion of run-in had some psychologic benefits for participants, with respect to demographic and depressive symptom characteristics, the TODAY cohort at baseline was similar to youth with type 2 diabetes in the population-based SEARCH for Diabetes in Youth Study (SEARCH) (8,9).

Older female adolescents reported more depressive symptoms in comparison with younger girls and with boys. A post hoc analysis revealed that only a small number of older youth ($n = 11$, all girls; 1.6%) scored at or above the usual cutoff score (≥ 29) for “severe” depressive symptoms. This pattern is similar to that of population samples where rates of depression are approximately the same for boys and girls during childhood, but diverge during adolescence when girls have depression rates two to three times higher than boys (10). Current results also are similar to those reported in the SEARCH study, which found that being female and older were risk factors for higher rates of depressive symptoms (11).

Depression is the most studied psychosocial factor in adolescents with type 1 diabetes, with prevalence rates for increased risk of elevated depressive symptoms reported between 14 and 33% (12,13). The risk for depression increases as

adolescents with type 1 diabetes develop into young adulthood, with Kovacs et al. (14) reporting that half of young adults with type 1 diabetes followed since adolescence developed a depressive or anxiety disorder.

There were no significant differences in depressive symptom or quality of life scores across racial/ethnic groups in either age-group. As expected, lower quality of life scores were significantly related to higher rates of depressive symptoms in both age-groups.

Because of the study design, at baseline the TODAY sample was relatively homogeneous, which may limit generalizability. Youth had low socioeconomic status and were predominately minorities, overweight/obese, and in adequate glycemic control at baseline. As the TODAY cohort becomes more heterogeneous over time with respect to glycemic control, BMI, and adherence, baseline depressive symptoms may become important predictors of subsequent treatment outcomes.

Although sample homogeneity is regarded as a study limitation, it is important to note that despite adequate glycemic control at baseline a large percentage of participants also had serious metabolic abnormalities present (3). Moreover, the sex, age, racial/ethnic groups, socioeconomic level, and depressive symptom characteristics of the TODAY cohort are similar to the those of the population-based SEARCH study (8,9), which suggests that the TODAY cohort is representative of the population of youth with type 2 diabetes in the U.S.

Table 1—Mean (SD) scores on the PedsQL, CDI (youth aged ≤ 15 years), and BDI (youth aged ≥ 16 years), and percent scoring at or above the cutoff score for “serious level of depressive symptoms” on each inventory and for combined CDI and BDI by racial/ethnic group and by sex

	PedsQL (all ages)		CDI (aged 10–15 years)			BDI (aged ≥ 16 years)			CDI ≥ 13 or BDI ≥ 14 (%)
	N	Mean (SD)	N	Mean (SD)	≥ 13 (%)	N	Mean (SD)	≥ 14 (%)	
Total	689	80.02 (12.04)	513	7.08 (6.36)	14.8	174	6.36 (7.55)	14.9	14.8
Race/ethnicity									
American Indian	42	78.23 (13.17)	28	6.50 (7.62)	14.3	15	7.87 (7.11)	13.3	14.0
Black non-Hispanic	217	79.58 (12.35)	166	7.02 (5.79)	15.1	50	6.02 (7.21)	14.0	14.8
Hispanic	282	80.54 (11.33)	213	7.38 (6.13)	15.0	68	7.40 (8.69)	20.6	16.4
White non-Hispanic	137	80.09 (12.44)	97	6.68 (7.41)	14.4	38	4.21 (5.29)	5.3	11.9
Asian non-Hispanic	11	81.25 (15.08)	9	7.22 (6.22)	11.1	3	8.00 (10.44)	33.3	16.7
P value*		0.63		0.77	0.99		0.17	0.20	0.68
Sex									
Female	450	79.49 (12.13)	351	7.19 (6.65)	15.1	101	8.27 (8.79)	22.8	16.8
Male	239	81.01 (11.83)	162	6.85 (5.68)	14.2	73	3.71 (4.18)	4.1	11.1
P value		0.11		0.58	0.79		<0.001	<0.001	0.044

*P values across racial/ethnic groups were computed without Asian non-Hispanic because of the small sample size.

Type 2 diabetes doubles the risk of depression in adults, with less than 25% receiving adequate depression treatment (15). Given the serious negative physical and psychosocial complications in depressed adults with type 2 diabetes (2,15), clinicians caring for older youth with type 2 diabetes should be familiar with mental health resources. Given the trajectory documented in adolescents with type 1 diabetes and high depressive symptoms for these symptoms to worsen as the older teen transitions into young adulthood, it is imperative to monitor and refer older adolescents with type 2 diabetes and depressive symptoms.

Acknowledgments—This work was completed with funding from the National Institute of Diabetes and Digestive and Kidney Diseases/ National Institutes of Health grants U01-DK-61212, U01-DK-61230, U01-DK-61239, U01-DK-61242, and U01-DK-61254; from the National Center for Research Resources General Clinical Research Centers Program grants M01-RR00036 (Washington University School of Medicine), M01-RR00043-45 (Children's Hospital Los Angeles), M01-RR00069 (University of Colorado Denver), M01-RR00084 (Children's Hospital of Pittsburgh), M01-RR01066 (Massachusetts General Hospital), M01-RR00125 (Yale University), and M01-RR14467 (University of Oklahoma Health Sciences Center); and from the National Center for Research Resources Clinical and Translational Science Awards grants UL1-RR024134 (Children's Hospital of Philadelphia), UL1-RR024139 (Yale University), UL1-RR024153 (Children's Hospital of Pittsburgh), UL1-RR024989 (Case Western Reserve University), UL1-RR024992 (Washington University), UL1-RR025758 (Massachusetts General Hospital), and UL1-RR025780 (University of Colorado Denver).

No potential conflicts of interest relevant to this article were reported.

The writing group takes responsibility for the content of this article: B.J.A. (chair), S.E., N.W.A., L.E.L.K., P.M.Y., P.M.T., S.E.T., S.V.M., T.L.C., and M.D.M. A list of the members of the

TODAY Study Group can be found in the Supplementary Data online.

B.J.A. and S.E. researched data, contributed to discussion, wrote the manuscript, and reviewed and edited the manuscript. N.W.A. contributed to discussion and reviewed and edited the manuscript. L.E.L.K. wrote, reviewed, and edited the manuscript. P.M.Y. researched data, contributed to discussion, and reviewed and edited the manuscript. S.J.L. researched data and contributed to discussion. P.M.T., S.E.T., and S.V.M. researched data, contributed to discussion, and reviewed and edited the manuscript. P.K. reviewed and edited the manuscript. T.L.C. researched data, contributed to discussion, and reviewed and edited the manuscript. M.D.M. contributed to discussion and reviewed and edited the manuscript.

Parts of this study were presented in abstract form at the 70th Scientific Sessions of the American Diabetes Association, Orlando, Florida, 25–29 June 2010.

The TODAY Study Group thanks the American Diabetes Association and the following companies for donations in support of the study's efforts: Becton, Dickinson and Company; Bristol-Myers Squibb; Eli Lilly and Company; GlaxoSmithKline; LifeScan, Inc.; Pfizer; and sanofi-aventis.

References

- Anderson RJ, Freedland KE, Clouse RE, Lustman PJ. The prevalence of comorbid depression in adults with diabetes: a meta-analysis. *Diabetes Care* 2001;24:1069–1078
- Lin EHB, Rutter CM, Katon W, et al. Depression and advanced complications of diabetes: a prospective cohort study. *Diabetes Care* 2010;33:264–269
- Copeland KC, Zeitler P, Geffner M, et al. Characteristics of adolescents and youth with recent-onset type 2 diabetes: the TODAY cohort at baseline. *J Clin Endocrinol Metab* 2011;96:159–167
- Kovacs M. *Children's Depression Inventory*. New York, Multi-Health Systems, 1992
- Beck AT, Steer RA, Brown GK. *Manual for the Beck Depression Inventory-II*. San Antonio, TX, Psychological Corporation, 1996
- Varni JW, Seid M, Rode CA. The PedsQL: measurement model for the Pediatric Quality of Life inventory. *Med Care* 1999;37:126–139
- Kessler RC, Avenevoli S, Ries Merikangas K. Mood disorders in children and adolescents: an epidemiologic perspective. *Biol Psychiatry* 2001;49:1002–1014
- Dabelea D, Bell RA, D'Agostino RB Jr, et al.; Writing Group for the SEARCH for Diabetes in Youth Study Group. Incidence of diabetes in youth in the United States. *JAMA* 2007;297:2716–2724
- Liese AD, D'Agostino RB Jr, Hamman RF, et al.; SEARCH for Diabetes in Youth Study Group. The burden of diabetes mellitus among US youth: prevalence estimates from the SEARCH for Diabetes in Youth Study. *Pediatrics* 2006;118:1510–1518
- Hankin BL, Abramson LY, Moffitt TE, Silva PA, McGee R, Angell KE. Development of depression from preadolescence to young adulthood: emerging gender differences in a 10-year longitudinal study. *J Abnorm Psychol* 1998;107:128–140
- Lawrence JM, Standiford DA, Loots B, et al.; SEARCH for Diabetes in Youth Study. Prevalence and correlates of depressed mood among youth with diabetes: the SEARCH for Diabetes in Youth study. *Pediatrics* 2006;117:1348–1358
- Grey M, Whittemore R, Tamborlane W. Depression in type 1 diabetes in children: natural history and correlates. *J Psychosom Res* 2002;53:907–911
- Hood KK, Huestis S, Maher A, Butler D, Volkering L, Laffel LM. Depressive symptoms in children and adolescents with type 1 diabetes: association with diabetes-specific characteristics. *Diabetes Care* 2006;29:1389–1391
- Kovacs M, Obrosky DS, Goldston D, Drash A. Major depressive disorder in youths with IDDM. A controlled prospective study of course and outcome. *Diabetes Care* 1997;20:45–51
- Rubin RR, Ciechanowski P, Egede LE, Lin EH, Lustman PJ. Recognizing and treating depression in patients with diabetes. *Curr Diab Rep* 2004;4:119–125